



Epileptiform activity triggers long-term plasticity of GABA(B) receptor signalling in the developing rat hippocampus.

Patrizia Tosetti, Nadine Ferrand, Isabelle Colin-Le Brun, Jean-Luc Gaïarsa

► To cite this version:

Patrizia Tosetti, Nadine Ferrand, Isabelle Colin-Le Brun, Jean-Luc Gaïarsa. Epileptiform activity triggers long-term plasticity of GABA(B) receptor signalling in the developing rat hippocampus.. The Journal of Physiology, 2005, 568 (Pt 3), pp.951-66. 10.1113/jphysiol.2005.094631 . inserm-00380221

HAL Id: inserm-00380221

<https://www.hal.inserm.fr/inserm-00380221>

Submitted on 13 May 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Epileptiform Activity Triggers Long-Term Plasticity of GABA_B Receptor Signalling in the
Developing Rat Hippocampus**

Abbreviated title: Long-Term Plasticity of GABA_B Receptor Signalling

P. Tosetti^{1,3}, N. Ferrand², I. Colin-Le Brun², and J.L. Gaiarsa²

¹ Institut de Génomique Fonctionnelle, CNRS UMR5203/INSERM U661/UM1/UM2, Montpellier, France.

² INMED, INSERM U29, Marseille, France

³**Correspondence should be addressed to:** Patrizia Tosetti, IGF, 144 rue de la Cardonille, 34094 Montpellier cedex 05, France.

Tel: 33 467 142 942

Fax: 33 467 542 432

E-mail: ptosetti@igf.cnrs.fr

Keywords: Desensitisation, GABA_B receptors, Epilepsy, Development.

Number of figures: 8

Number of tables: 1

Number of pages: 35

ABSTRACT

GABA_B receptor (GABA_BR)-mediated presynaptic inhibition regulates neurotransmitter release from synaptic terminals. In the neonatal hippocampus, GABA_BR activation reduces GABA release and terminates spontaneous network discharges called Giant Depolarising Potentials (GDPs). Blocking GABA_BRs transforms GDPs into longer epileptiform discharges. Thus, GABA_BR-mediated presynaptic inhibition of GABA release (GABA auto-inhibition) controls both spontaneous network activity and excitability in the developing hippocampus.

Here we show that extensive release of endogenous GABA during epileptiform activity impairs GABA auto-inhibition, but not GABA_BR-mediated inhibition of glutamate release, leading to hyperexcitability of the neonatal hippocampal network. Paired-pulse depression of GABA release (PPD) and heterosynaptic depression of glutamate release were used to monitor the efficacy of presynaptic GABA_BRs-mediated inhibition in slices. PPD, but not heterosynaptic depression, was dramatically reduced after potassium (K)-induced ictal-like discharges (ILDs), suggesting a selective impairment of GABA_BR-dependent presynaptic inhibition of GABAergic terminals. Impairing GABA auto-inhibition induced a 44% increase in GDP width and the appearance of pathological network discharges. Preventing GABA-induced activation of GABA_BRs during ILDs avoided PPD loss and most modifications of the network activity. In contrast, a partial block of GABA_BRs induced network discharges strikingly similar to those observed after K-driven ILDs. Finally, neither loss of GABA auto-inhibition nor network hyperexcitability could be observed following synchronous release of endogenous GABA in physiological conditions (during GDPs at 1 Hz). Thus, epileptiform activity was instrumental to impair GABA_BR-dependent presynaptic inhibition of GABAergic terminals.

In conclusion, our results indicate that endogenous GABA released during epileptiform activity can reduce GABA auto-inhibition and trigger pathological network discharges in the newborn rat hippocampus. Such functional impairment may play a role in acute post-seizure plasticity.

INTRODUCTION

The occurrence of seizures during development strongly increases the probability of developing chronic epilepsy at adult age (Holmes & Ben-Ari, 1998). This suggests that plastic changes occurring during and after seizures underlay subsequent remodelling. Ictal discharges are stereotyped epileptic events involving the synchronous activation of thousands of neurons in the cortex and hippocampus. Long-term changes associated with ictal activity have been thoroughly characterised in the developing brain, and include neuronal hyperexcitability (Villeneuve *et al.*, 2000; Bender *et al.*, 2003), glutamatergic axonal sprouting (Bender *et al.*, 2003; Holmes *et al.*, 1998) and modifications of ionotropic γ -aminobutyric acid (GABA)-receptor mediated synaptic transmission (Chen *et al.*, 1999; Khalilov *et al.*, 2003). Impaired GABA_B receptor (GABA_BR)-mediated signalling has been observed after epileptic activity in the adult hippocampus (Mangan & Lothman, 1996; Haas *et al.*, 1996; Chandler *et al.*, 2003; Wu & Leung, 1997). In the developing hippocampus, however, it is not clear whether GABA_BR-mediated signalling is functional after seizures.

GABA_BRs play a crucial role in controlling the physiological activity of the immature hippocampal network. GABA_A currents, which mediate most of the inhibitory drive in the adult, are excitatory during the first postnatal week of life (Cherubini *et al.*, 1991). Consequently, the synchronous activation of hippocampal interneurons results in excitatory network discharges called Giant Depolarising Potentials (GDPs) (Ben-Ari *et al.*, 1989). In these peculiar conditions, GABA_BR activation is essential to limit network excitability. Endogenous GABA released during GDPs activates GABA_BRs, thus reducing further GABA release from interneurons and leading to GDP termination (McLean *et al.*, 1996). Consistent with this, the pharmacological block of GABA_BRs progressively increases GDP width, transforming GDPs into epileptiform discharges (McLean *et al.*, 1996; Tosetti *et al.*, 2004). Similar effects can also be observed after the functional impairment of GABA_BR-mediated presynaptic inhibition of GABA release (GABA auto-inhibition) by a prolonged application of baclofen, a selective GABA_BR agonist (Tosetti *et al.*, 2004). This observation raises the interesting possibility that a sustained release of endogenous GABA may trigger a functional loss of GABA auto-inhibition, leading to hyperexcitability and pathological network activity. Ictal discharges,

during which both GABA and glutamate are continuously released for several seconds (Velazquez & Carlen, 1999; Kohling *et al.*, 2000; Rutecki *et al.*, 1985), are one possible context for such loss to occur.

Here we report that extensive endogenous GABA release during ictal-like discharges impairs GABA auto-inhibition in the newborn rat hippocampus. Such impairment contributes to the network hyperexcitability observed following the epileptiform discharges. These data thus reveal a novel role of GABA_BR-mediated signalling in acute post-seizure plasticity of the developing hippocampus.

METHODS

Tissue preparation

Brains were removed from anaesthetised (350 mg kg^{-1} chloral hydrate intraperitoneally) male Wistar rats (postnatal day 2-5) and immersed into ice cold ($2-4^{\circ}\text{C}$) artificial cerebrospinal fluid (ACSF) of the following composition (mM): NaCl, 126; KCl, 3.5; CaCl_2 , 2; MgCl_2 , 1.3; NaH_2PO_4 , 1.2; NaHCO_3 , 25; and glucose, 11; pH 7.4, when equilibrated with 95% O_2 and 5% CO_2 . All animal experiments were carried out in accordance with the European Communities Council Directive of 24 November 1986 (86/609/EEC). Intact hippocampal formations (IHF) were dissected out as previously described (Khalilov *et al.*, 1997), and incubated in oxygenated ACSF at room temperature. When needed, $600 \mu\text{m}$ slices were prepared from IHFs using a McIlwain tissue chopper. Both intact IHFs and slices were equilibrated at room temperature in oxygenated ACSF for at least 60 min prior to electrophysiological recordings.

Recording solutions

ACSF, the extracellular recording solution, was prepared from a 10 X stock solution immediately before the experiment, equilibrated with 95% O_2 and 5% CO_2 , heated at 34°C , and delivered to the recording chamber via a pressurised system at a flow rate of 3 ml min^{-1} (slices), or 10 ml min^{-1} (IHF).

For whole-cell recordings from hippocampal slices, intracellular solution contained (in mM): 150 CsCl₂, 2 MgCl₂, 0.1 CaCl₂, 1 EGTA, 2.5 Na₂ATP, 0.4 Na₂-GTP, 1.5 Mg-ATP, 10 HEPES, pH=7.25, 2QX314, osmolarity=275 mOsmol. In these ionic conditions, both ionotropic GABAergic and glutamatergic currents reversed around 0 mV. For patch-clamp recordings of the IHF, the intracellular solution contained (in mM): 120 CsGluconate, 13.4 CsCl, 10 HEPES, 1.1 EGTA, 0.1 CaCl₂, 0.4 Na₂GTP, 4 MgATP, pH=7.4. In these ionic conditions, GABA_A currents reversed around -50 mV , while ionotropic glutamatergic currents reversed around $+10 \text{ mV}$. For all patch-clamp experiments, 5,6-tetramethylrhodamine biocytin (rhodamine) was dissolved (0.5-1%) in the internal solution to

allow post-hoc visualisation of the recorded neurons. For extracellular field recordings, pipettes were filled with ACSF.

Electrophysiological recordings

CA3 pyramidal neurons were recorded from either hippocampal slices or IHFs (Khalilov *et al.*, 1997). Spontaneous and evoked synaptic activities were acquired with an Axopatch 2B patch-clamp amplifier (Axon Instruments, USA), filtered at 10 kHz, digitised using a Digidata 1200 A/D interface (Axon Instruments, USA) and stored on a Pentium II 450 MHz PC running Axoscope 8.1 (Axon Instruments, USA) for off-line analysis. Field potentials were recorded using a DAM80 Amplifier (World Precision Instruments, USA) using a 1 to 3 Hz band pass filter. Capacitance, input and series resistances were measured online with Acquis Software (Biologic, France). Patch-clamp pipettes had a resistance of 3 to 5 M Ω , once filled with the internal recording solution. Series resistances ranged from 15 to 30 M Ω . Cells showing series resistances > 30 M Ω were discarded. Field recordings were performed with glass micropipettes of 10-20 M Ω resistance. To reserve the viability of the IHF preparation and avoid ischemic artefacts, each IHF was recorded for no more than 2 hrs.

Impairment of GABA auto-inhibition

In both hippocampal slices and IHFs, functional impairment of GABA auto-inhibition was induced via ictal-like discharge (ILD)-promoted sustained release of endogenous GABA (Fig 1A). ILDs were triggered by the bath application (9.1 ± 0.9 min, range 8-10 min) of 8.5 to 10 mM K⁺. When either 500 μ M CGP35348 or 10 μ M CNQX + 40 μ M APV were co-applied with high K, the cumulative discharge time was monitored to guarantee GABA release consistency among pharmacologically distinct induction phases. In high K, ILDs represented $41 \pm 3\%$ of the overall induction duration. In high K + CGP35348, the ILD contribution was not significantly different ($39 \pm 2\%$, $P > 0.1$), although the presence of CGP35348 slightly shortened the ILD duration (high K: 90 ± 5 s, $n=30$; high K + CGP35348: 76 ± 3 s, $n=12$, $P < 0.05$) and increased their frequency (high K: 4.1 ± 0.1 mHz; high K + CGP35348: 5.6 ± 0.2 mHz; $P < 0.01$). In high K + CNQX + APV, ILDs were not present, but GABA

was still released via short, frequent bursts that constituted $63 \pm 4\%$ of the induction phase. When ILDs were induced by 500 μM CGP35348 alone, they represented $44 \pm 5\%$ of the induction phase.

Data analysis

All recordings were visualised using Axoclamp 8.1 software (Axon Instruments, USA), and analysed using the Acquis (Biologic, France), Minianalysis (Synaptosoft, USA) and Igor software (Wavemetrics, USA).

Paired pulse depression (PPD) of GABAergic synapses impinging onto CA3 pyramidal neurons was measured using pairs of identical stimuli (4-15 V, 0.03 ms each) at 250-350 ms interval, delivered at a frequency of 0.03 Hz with a bipolar tungsten electrode placed in the CA3 *stratum radiatum*. The resulting pairs of postsynaptic GABA_A currents (GABA_A-PSCs) were isolated in the presence of 10 μM CNQX and 40 μM D-APV. PPD was measured from the amplitude difference between the 1st and 2nd GABA_A-PSCs, normalised to the amplitude of the 1st GABA_A-PSC. Average PPD values were calculated from 15 paired stimulations delivered at 0.03 Hz starting either 15 min before (control PPD) or 5 min after the different pharmacological treatments. For baclofen experiments, a 10 min ACSF wash was introduced between the end of the baclofen treatment and the restarting of the PPD stimulations, to allow a thorough washout of the drug. Cells showing differences in the amplitude of the 1st GABA_A-PSC before and after treatment were discarded.

Heterosynaptic depression of glutamatergic synapses impinging on CA3 pyramidal neurons was measured by stimulating two independent glutamatergic pathways via two stimulating electrodes positioned in the *stratum radiatum* on opposite sides of the recording electrode. The electrode position and stimulus intensity were optimized for maximal field EPSP (fEPSP) amplitude. Stimulated pathways were tested for independence by verifying the absence of cross-facilitation. To induce heterosynaptic depression, a single stimulus was given to the 1st pathway (unconditioned fEPSP). After 30 sec, an identical stimulus was given to the same pathway, immediately (300 ms) preceded by a conditioning train (10 stimuli at 50 Hz) to the 2nd pathway. This originated a conditioned fEPSP. The magnitude of heterosynaptic depression was calculated from the amplitude difference between the

unconditioned and conditioned fEPSPs, normalised to the amplitude of the unconditioned fEPSP. All recordings were performed in the presence of 20 μ M tiagabine, a selective blocker of GABA neuronal transporter 1.

To characterise the physiological activity of the hippocampal network, we measured the amplitude and duration of the Giant Depolarising Potentials (GDPs) that are spontaneously present in the untreated IHF. GDP duration was measured at 20% of GDP amplitude. To characterise the pathological network activity following the different pharmacological treatment, we measured the amplitude, duration and shape of abnormal events recorded extracellularly from the IHF. Event shape was quantified by the number of phases within each event, where 1 phase is defined as 1 change in the slope of the recorded trace. Amplitude and duration of abnormal events were normalised to the respective GDP values from the same cell. Frequency distributions of event amplitude, duration and shape were built for individual cells. Those were then averaged to generate mean frequency histograms for each parameter.

Ionotropic GABA and glutamatergic currents underlying network events were isolated by means of their distinct reversal potentials while using CsGluconate internal solution. GABA and glutamate charge transfers for each event were then calculated from the areas of the GABAergic and glutamatergic currents underlying the event itself.

Statistical analysis was performed with Kolmogorov-Smirnov (K-S), ANOVA or Student's T tests, as appropriate. All values were expressed as mean \pm standard error. Differences were considered significant when $P < 0.05$.

Post-hoc morphological characterisation of recorded cells

After recordings, slices were fixated overnight at 4°C in 0.1 M phosphate buffer saline (PBS) containing 4% paraformaldehyde and 0.9% NaCl. Slices were then rinsed in PBS, mounted on gelatin-coated slides with an aqueous mounting medium (Gel Mount, Biomedex), and coverslipped. Rhodamine-filled cells were identified with an Olympus confocal microscope (Fluoview BX50WI) using a helium/neon laser as the excitation source (543 nm) and an emission filter band pass (560nm).

Images were taken with a digital camera (ORCA-ER, Hamamatsu). Cells were classified as either pyramidal neurons or interneurons on the basis of their morphology and axonal projections. Electrophysiological data from interneurons were discarded.

Drug delivery

Drugs were diluted from stock solutions into freshly prepared ACSF immediately before the experiment. Drugs used in this study were: Baclofen (Sigma), 6-cyano-7-nitroquinoxaline-2,3-dione (CNQX; Tocris), D(-)-2-amino-5-phosphovaleric acid (D-AP5; Tocris), P-3-aminopropyl-P-diethoxymethyl phosphoric acid (CGP35348, Novartis, Basel), tiagabine (Novo Nordisk, Denmark).

RESULTS

Impairing GABA_B receptor-mediated presynaptic inhibition

Our working hypothesis was that, during epileptiform activity, the sustained exposure of GABA_B receptors (GABA_BRs) to elevated quantities of endogenous GABA could impair GABA_BR-mediated presynaptic inhibition of GABA release (GABA auto-inhibition). Such functional loss would then increase hyperexcitability and induce spontaneous pathological discharges in the newborn rat hippocampus.

To test this idea, we first investigated whether ictal-like discharges (ILDs) released enough GABA to impair GABA auto-inhibition. ILDs were induced by application of 8.5 to 10 mM K (9.1 ± 0.9 min, range 8-10 min) to either hippocampal slices or the intact hippocampal formation (IHF). We chose this acute model of epilepsy because elevation of $[K]_o$ occurs during seizures and contributes to their generation (Freund *et al.*, 1990; Jensen & Yaari, 1997). High K induced interictal-like activity followed by ILDs lasting 10-20 sec in slices, and 40-120 sec in the IHF (Fig. 1A). ILDs always started with tonic oscillations at 10-20 Hz and ended with clonic bursts occurring at 1-2 Hz (Fig 1A). In most cases, an intermediate phase of tonico-clonic bursts at variable frequency could also be observed (Fig 1A). Paired extracellular field recordings from the *stratum radiatum*, and whole cell recordings from CA3 pyramidal neurons demonstrated that GABA was continuously released over tens of seconds before and during ILDs in the IHF (Fig. 1A, *upper trace*). These are therefore ideal conditions for inducing a functional impairment of GABA auto-inhibition, since GABA_B receptors are exposed to a sustained and intense activation during each discharge.

Paired-pulse depression and heterosynaptic depression as indexes of efficacy of GABA_BR-mediated presynaptic inhibition

Paired-pulse depression at GABAergic synapses (PPD) (Davies *et al.*, 1990) and heterosynaptic depression at glutamatergic synapses (Chandler *et al.*, 2003) were used to monitor the efficacy of presynaptic GABA_BR-mediated inhibition in hippocampal slices.

Both PPD and heterosynaptic depression depended upon GABA_B receptor activation, as they were severely reduced by the selective GABA_BR antagonist, CGP35348. As shown in Fig 1B, average PPD was reduced from $35\pm6\%$ (n=6) in control conditions to $7\pm2\%$ in the presence of 500 μ M CGP35348 (n=6, $P<0.005$). Similarly, heterosynaptic depression was reduced from $35\pm3\%$ to $12\pm3\%$ in the presence of the GABA_BR antagonist (Fig. 1C). Both PPD and heterosynaptic depression were fully restored following CGP35348 washout. These results indicate that both PPD and heterosynaptic depression rely upon activation of presynaptic GABA_B receptors in our experimental conditions. The functional impairment of presynaptic GABA_BR-mediated inhibition should therefore result in a reduction of both PPD and heterosynaptic depression.

In a previous work (Tosetti *et al.*, 2004), we showed that, among the different GABA_B receptor-mediated responses, only the presynaptic inhibition of GABA release was impaired after a prolonged application of baclofen. We confirmed this result by measuring PPD and heterosynaptic depression before and after a 5 min application of 100 μ M baclofen, a GABA_BR agonist. While in the bath, baclofen induced a decrease in the probability of GABA release, resulting in a reduction of the 1st pulse amplitude (Fig. 2A, left panel), and a switch from depression to facilitation (control: $26\pm5\%$; during baclofen: $-28\pm16\%$; n=6, $P<0.005$) (Fig. 2A *control, during baclofen*). Interestingly, PPD did not recover once baclofen was washed off, and it was still absent ($-1\pm2\%$) 20 min after removal of the agonist (n=6, $P<0.005$) (Fig 2A, *after baclofen*). This could not be accounted for by an incomplete baclofen washout, since the amplitude of the first pulse recovered to pre-baclofen values (Fig. 2A, left panel) and the facilitation disappeared (Fig. 2A, right panel). These results therefore suggest a baclofen-induced loss of GABA auto-inhibition.

In contrast, heterosynaptic depression recovered to control values after baclofen washout (Fig. 2B, *wash*). Heterosynaptic depression was $36\pm2\%$ in control conditions, $4\pm4\%$ during, and $51\pm9\%$ after baclofen application (n=6, Fig. 2B). Baclofen therefore did not impair GABA_BRs on glutamate fibers.

In conclusion, these results confirm that sustained activation of GABA_BRs results in a selective decrease in the efficacy of presynaptic GABA_BR-mediated inhibition of GABA release. These results also show that PPD and heterosynaptic depression can be used as indexes of functional impairment of

GABA_BR-mediated presynaptic inhibition in conditions of conserved GABA uptake and release probability (Roepstorff & Lambert, 1994).

ILD-induced release of GABA impairs GABA auto-inhibition

We next investigated whether endogenous GABA released during ILDs could impair GABA auto-inhibition in hippocampal slices. 6 ILDs were induced via an 8-10 min application of 10 mM K (induction phase), as described above. We measured PPD and heterosynaptic depression before and following the induction phase, and found that only PPD was severely reduced with respect to control (Fig. 3). Average PPD was decreased from $34\pm4\%$ in control conditions to $8\pm3\%$ after K treatment ($n=8$; $P<0.001$; Fig. 3A). It should be noted that the amount of residual PPD after K treatment is consistent with the amount of GABA_BR-independent PPD measured in the presence of CGP35348. In contrast, average heterosynaptic depression did not significantly differ before (33 ± 3 ; Fig. 3B, *Control*) and after the induction phase ($43\pm9\%$, $n=7$; Fig. 3B, *After high K*). Thus, in agreement with the results of the baclofen treatment, high K treatment selectively affected GABA auto-inhibition.

Although a decreased PPD is consistent with the functional impairment of presynaptic GABA_BR signalling, other factors may contribute to it. In fact, PPD depends also on GABA release probability and uptake (Roepstorff & Lambert, 1994). A decreased GABA release probability was unlikely since the amplitude of the GABA_A postsynaptic current evoked by the 1st stimulus (1st GABA-PSC, see stimulating protocol in Fig 1B) was preserved before and after high K treatment (Fig. 3A, left panel). In agreement with this, the coefficient of variation of the 1st GABA-PSC amplitude (CV_A) did not significantly differ before (0.13 ± 0.02) and after high K treatment (0.14 ± 0.03 ; $n=8$, $P>0.3$). Thus, the PPD loss cannot be accounted for by a decreased probability of GABA release. In control conditions, the CV_A of the 2nd GABA-PSC was much higher than that of the 1st (0.41 ± 0.1 , $P<0.01$), in agreement with a reduced probability of release following GABA_BR activation. However, such value decreased significantly after high K treatment (0.18 ± 0.04 , $P<0.01$), as expected after a loss of GABA auto-inhibition.

We next tested the possibility that PPD loss was due to ILDs reducing neuronal GABA uptake. PPD was measured in the presence of 20 μ M tiagabine, a selective blocker of GABA neuronal transporter 1 (Andersen *et al.*, 1993), both before and after ILDs. PPD in tiagabine was $48\pm 2\%$ before and $8\pm 4\%$ after high K treatment ($n=9$; $P<0.001$; data not shown). ILDs therefore significantly reduced PPD even in the absence of functional neuronal GABA transporters, ruling out an impaired neuronal GABA uptake as the cause for PPD loss.

Since neither GABA uptake nor the probability of GABA release were affected by ILDs, functional impairment of presynaptic GABA_BR signalling is the most likely cause of high K-induced PPD loss. To confirm such conclusion, we tested whether PPD loss required the activation of GABA_BR by co-applying 500 μ M CGP35348 during the high K induction phase. Interestingly, no PPD loss was observed after the high K+CGP35348 treatment (Fig. 4A). Average PPD was $27\pm 2\%$, not significantly different from that measured in control conditions, $27\pm 4\%$ ($P>0.6$; $n=9$). Thus, these results show that activation of GABA_BRs by endogenous GABA is required to induce PPD loss.

We next investigated whether the pattern of GABA_BR activation played an important role in the impairment of GABA auto-inhibition. Co-applying 40 μ M APV + 10 μ M CNQX during the high K induction phase eliminated ILDs, leaving only shorter (500 ms) GABAergic bursts at 0.8 Hz (Fig 4B). Such discharges, which are reminiscent of Giant Depolarising Potentials (GDPs), release sufficient endogenous GABA to activate GABA_BRs (McLean *et al.*, 1996), but are probably too short to functionally impair GABA_BR-mediated signalling. Indeed, GABA_BRs are functional in the presence of GDPs in physiological conditions (McLean *et al.*, 1996). Interestingly, PPD was not significantly different before ($32\pm 7\%$) and after K+CNQX+APV ($42\pm 12\%$; $n=6$; $P>0.05$; Fig 4C). Thus, the intermittent activation of GABA_BRs does not have any effect on PPD. These results show that it is not GABA_BR activation *per se*, but sustained GABA_BR activation that is responsible for decreasing PPD.

In conclusion, our results show that endogenous GABA released during epileptiform activity can induce the functional impairment of GABA auto-inhibition in the newborn hippocampus. This effect requires a sustained GABA release. Intermittent, short GABAergic discharges mimicking physiological GDPs are ineffective.

ILD-induced functional impairment of GABA auto-inhibition modifies network activity in hippocampal slices

Our next goal was to investigate the acute and long-term consequences of a reduced GABA auto-inhibition on the immature hippocampal network. GDPs, the giant depolarising potentials, constitute the hallmark of early hippocampal network activity (Ben-Ari *et al.*, 1997). In hippocampal slices, GDP duration was increased by $44\pm 8\%$ following 6 high K-induced ILDs (Fig. 5, *after high K*). Average GDP width (measured at 20% of GDP amplitude) increased from 300 ± 14 ms in control conditions to 432 ± 20 ms after ILDs ($n=6$; $P<0.01$). Such increase is consistent with a functional loss of GABA_BR-mediated presynaptic inhibition, since previous work demonstrated that GABA_BR activation is required for GDP termination (Tosetti *et al.*, 2004).

We next confirmed that the increased GDP duration depended on GABA_BR activation by applying 500 μ M CGP35348 during the high K induction phase. Such treatment indeed prevented the increase in GDP width (control: 295 ± 17 ms; after high K+CGP35348: 326 ± 23 ms; $n=6$; $P>0.2$) (Fig. 5, *after high K+CGP35348*). These results indicate that GABA_BR activation is required to increase GDP duration, and support the idea that ILD-induced impairment of GABA auto-inhibition accounts for the majority of the observed increase in GDP width. An alternative approach to the same question was to induce ILDs via the application of 500 μ M CGP35348 alone (without elevating extracellular K). In fact, by blocking GABA_B receptors, saturating concentrations of CGP35348 transformed spontaneous GDPs into ILDs (McLean *et al.*, 1996; Tosetti *et al.*, 2004). This strategy had the dual advantage of generating a pattern of GABA release similar to that induced by high K (see Methods) while preserving GABA_B receptor function after the induction phase. CGP35348-induced ILDs did not affect GDP duration, which passed from 289 ± 27 ms to 316 ± 30 ms after treatment ($P>0.1$; Fig 5, *after CGP35348*).

We finally investigated the effects on GDP duration of a pattern of GABA release unable to impair GABA auto-inhibition. The short, frequent GABAergic bursts generated by the co-application of high K and 10 μ M CNQX+40 μ M APV (see previous chapter and Fig. 4B) did not significantly increase

GDP width, which was 312 ± 49 ms in control conditions and 335 ± 44 ms after treatment ($n=8$; $P>0.5$). Thus, GABA_BR activation by endogenous GABA could not *per se* increase GDP width, unless it was long enough to impair GABA auto-inhibition. It is therefore the ILD-induced decrease of GABA auto-inhibition that is responsible for increasing GDP duration.

To further support this conclusion, we tested the effect on GDP width of a 5 min application of 100 μ M baclofen, a treatment known to impair GABA auto-inhibition (Fig. 2) (Tosetti *et al.*, 2004). As expected, GDP duration after baclofen treatment was significantly increased by $50 \pm 8\%$, from 275 ± 8 ms to 410 ± 18 ms ($n=8$, $P<0.001$). Interestingly, this increase is not significantly different from that measured after high K treatment ($P>0.4$; Fig 5A).

In conclusion, our results indicate that ILD-induced GABA release can increase GDP width and modify the spontaneous network activity by impairing GABA_BR-mediated GABA auto-inhibition in immature hippocampal slices.

ILD-induced impairment of GABA auto-inhibition modifies network activity in the intact hippocampal formation

Since hippocampal slices contain only a fraction of the hippocampal network, they may not ideal to evaluate network activity. We therefore decided to further investigate the consequence of K-induced ILDs using the intact hippocampal formation (IHF), a preparation that preserves the hippocampal circuitry (Khalilov *et al.*, 1997).

Network events occurring before and after high K-induced ILDs were monitored using extracellular field recordings of CA3 *stratum radiatum*. Events were then classified according to their amplitude, duration and shape (quantified as number of phases, see Methods). We defined as abnormal all events that significantly differed from GDPs in at least 2 out of the 3 above parameters.

In control conditions, GDPs constituted the totality of network events ($n=24$; Fig. 6A, *Control*) and could be described as stereotyped triphasic waveforms of 21 ± 2 μ V amplitude (range 8-43 μ V; $n=24$), and 484 ± 48 ms duration (range 131-692 ms). Following ILDs, however, GDPs accounted for only $26 \pm 4\%$ of total events ($n=24$; Fig. 6A, *after high K*). The remaining $74 \pm 4\%$ of network activity was

composed of abnormal events that were never recorded in control conditions. They could be divided into three major classes: short oscillations, long oscillations, and interictal-like events (Fig. 6B). Short oscillations were characterised by relatively small amplitude ($172 \pm 19\%$ of average GDP amplitude; range 35-405%; $n=25$), intermediate duration ($424 \pm 53\%$ of average GDP duration; range 155-1161%) and intermediate number of phases (8.0 ± 0.8 phases; range 2-15; GDPs=3 phases). Long oscillations not only displayed longer duration ($3907 \pm 797\%$ of average GDP duration, range 1097-9658%; $n=10$), but also larger amplitude ($942 \pm 334\%$ of average GDP amplitude, range 320-3167%), and always more than 15 phases. Interictal-like events were mostly triphasic (average= 3.2 ± 0.2 , range 2-5; $n=14$), with intermediate amplitude ($478 \pm 94\%$ of average GDP amplitude, range 103-1444%), and relatively short duration ($196 \pm 19\%$ of average GDP duration, range 35-438%). All three types of abnormal events were spontaneously present in the IHF up to at least 1 hr 30 min after the induction phase.

Simultaneous patch clamp and field recordings of the IHF were used to characterise the conductances underlying both GDPs and abnormal events (Fig. 6B). As previously reported, control GDPs were mainly driven by GABAergic currents (Ben-Ari *et al.*, 1989; Khazipov *et al.*, 1997). Their average GABA/glutamate charge transfer ratio was 16 ± 3 ($n=10$), indicating that ionotropic glutamatergic currents accounted for only 6.25% of total GDP current. Abnormal waveforms, however, were driven by sizeable amounts of both GABAergic and glutamatergic currents. Their average GABA/glutamate charge transfer ratio was 2.9 ± 0.4 ($n=11$), indicating that ionotropic glutamatergic currents represented approximately 34% of the total event current. Thus, GDPs and abnormal waveforms constitute distinct events.

In addition to releasing endogenous GABA and impairing the GABA_BR-dependent response, high K-induced ILDs also released large quantities of glutamate (Rutecki *et al.*, 1985). Strong glutamate release could induce several forms of plasticity that could contribute to the observed changes in network activity (Ben-Ari & Gho, 1988). To identify which type of abnormal event depended upon the functional loss of GABA auto-inhibition, we prevented GABA_BR activation during sustained GABA release with the application of 500 μ M CGP35348. This resulted in the recovery of most of the spontaneous GDP activity ($70 \pm 8\%$ of total events, $n=12$; Fig. 6A, *after high K + CGP*). The remaining

30% of events consisted exclusively of interictal-like waveforms. These results suggest that a decreased GABA auto-inhibition is required for the appearance of both long and short oscillations, but not interictal-like events in the immature hippocampus.

Fig. 7 shows frequency distributions of abnormal event properties (duration, shape, and amplitude) measured after high K alone or high K+500 μ M CGP3534 (n=24; for histogram values see Table 1). Superposed rectangles highlight properties whose frequency did not significantly differ in the presence or absence of GABA_BR activation. Interestingly, the duration, shape, and amplitude values that were independent of GABA_BR activation well matched the properties of interictal-like events. These results confirm that preserving GABA_BR-mediated signalling prevented the appearance of long and short oscillations.

If long and short oscillations are consequences of an impaired GABA auto-inhibition, treatments leading to a partial loss of function of GABA_BRs should trigger similar discharges. To test such hypothesis, we reduced GABA_BR-mediated signalling by applying non-saturating doses of CGP35348. Indeed, long and short oscillations strikingly similar to those recorded after high K-induced ILD could be recorded during the application of 100 μ M CGP35348 (Fig. 8A). CGP35348-driven events were mediated by both GABAergic and glutamatergic ionotropic currents (Fig. 8A). Their GABA/glutamate charge transfer ratio was 3.4 ± 0.8 (n=6), a value not significantly different from that of K-induced oscillations. Even more striking, the type and frequency of events recorded in CGP35348 and after high K-induced ILDs closely matched. After high K treatment, GDPs accounted for $27 \pm 3\%$ of total events, short oscillations for $57 \pm 5\%$, and long oscillations for $5 \pm 2\%$ (Fig. 8B). Similarly, during 100 μ M CGP35348, $32 \pm 3\%$ of the events were GDPs, $63 \pm 3\%$ were short oscillations, and $5 \pm 1\%$ were long oscillations (Fig. 8B). Interestingly, interictal-like events were not present during CGP application (Fig. 8B), in agreement with the idea of these events being a non-specific effect of ILDs rather than a consequence of impairing GABA auto-inhibition.

In conclusion, the partial block of GABA_B receptors induces the appearance of abnormal events that are qualitatively and quantitatively similar to those recorded after high K-induced ILDs. These

results indicate that ILD-induced GABA release impairs GABA auto-inhibition and triggers the onset of abnormal short and long oscillations in the immature hippocampus.

DISCUSSION

Our study focused on the functional consequences of acute epileptiform activity (ictal-like discharges, ILDs) on presynaptic GABA_BR-mediated signalling and network activity in the developing rat hippocampus. The central results were: 1) endogenous GABA release could impair GABA_BR-mediated presynaptic inhibition of GABAergic terminals (GABA auto-inhibition); 2) ILDs, but not physiological GDPs, provided the sustained GABA release required to induce such impairment; 3) the ILD-dependent loss of GABA auto-inhibition modified the spontaneous network activity and induced the onset of abnormal discharges, possibly contributing to post-seizure plasticity of the developing hippocampus.

Endogenous GABA impairs GABA auto-inhibition

We have demonstrated that high K-induced epileptiform activity can significantly reduce paired pulse depression (PPD) in the CA3 region of the developing rat hippocampus. PPD loss occurred after as few as 6 ILDs. Interestingly, a similar decrease in PPD could be observed in the CA1 region and in dentate gyrus granule cells after kindling (Wu & Leung, 1997; Buhl *et al.*, 1996).

How does such functional loss occur? Our data indicate that a sustained release of endogenous GABA during ILDs induces a PPD loss that is the consequence of an impaired GABA auto-inhibition. To support this hypothesis, we showed that activation of GABA_BRs during high K-induced ILDs was required to induce the PPD loss since: i) the application of the specific GABA_BR antagonist CGP 35348 during the induction phase prevented it and ii) a 5 min baclofen application induced a similar PPD loss. We also proved that GABA_BR activation needed to be long-lasting in order to induce the PPD loss. In fact, PPD was not reduced when the pattern of GABA_BR activation was changed from sustained (during ILDs) to intermittent (during GDP-like activity). Finally, we showed that other mechanisms potentially able to induce PPD loss, like a decreased probability of GABA release or an increased GABA uptake (Roepstorff & Lambert, 1994), were not involved. Altogether, these

observations indicate that, in the newborn hippocampus, GABA released during ILDs impairs presynaptic GABA_BR-mediated inhibition, leading to PPD loss.

The mechanism leading to the functional impairment of presynaptic GABA_BR-mediated inhibition is presently unknown. The simplest explanation is possibly that the release of endogenous GABA is sufficient to induce a conformational change of the GABA_B receptor, leading to a desensitized state insensitive to the agonist. For example, the impaired GABA auto-inhibition could result from uncoupling to the effectors via agonist induced dephosphorylation of the GABA_B receptors (Couve *et al.*, 2002). Alternatively, it could result from phosphorylation by G-protein receptor kinases (GRKs) which, through the binding of β -arrestins, target the G-protein-coupled receptors for endocytosis (Tsao & von Zastrow, 2000; Carman & Benovic, 1998; Ferguson *et al.*, 1998). Independently of the mechanism involved, the present study provides the first demonstration that endogenous GABA can impair GABA_BR-mediated inhibition and reveals one functional context where this occurs: during epileptiform activity.

Functional consequences of ILD-induced impairment of GABA auto-inhibition

We have shown that the ILD-induced functional loss of GABA auto-inhibition increased the duration of physiological Giant Depolarising Potentials (GDPs) and triggered pathological network oscillations in the intact hippocampus. In fact, procedures that prevented such impairment (co-application of CGP35348 or CNQX+APV during the high K induction phase) also prevented alterations of network activity. Furthermore, procedures that induced (bath application of baclofen) or simply mimicked impairment (non-saturating concentrations of CGP35348) triggered similar abnormal oscillations and increased GDP duration.

These results are consistent with previous studies reporting the fundamental role of GABA_BRs in preventing excessive hippocampal excitability during development. In fact, during the first postnatal week, the synchronous activation of GABAergic interneurons triggers depolarising GABA_AR-mediated currents and contributes to spontaneous network discharges called Giant Depolarising Potentials (GDPs) (Cherubini *et al.*, 1991). However, once released, GABA also acts on presynaptic

GABA_BRs, limiting further GABA release and effectively leading to GDP termination (McLean *et al.*, 1996). Impairing GABA_BR function at this age prevents GDP termination, transforming GDPs into long epileptiform discharges (Tosetti *et al.*, 2004). In agreement with these findings, our present data show that ILDs impair GABA auto-inhibition and decrease the inhibitory control exerted by endogenous GABA on the network activity. As a direct consequence, the hippocampal network develops a situation of hyperexcitability, revealed by the presence of longer GDPs, and abnormal long and short oscillations.

In the hippocampus, GABA_BRs are ubiquitously located at both postsynaptic and presynaptic sites on both GABAergic and glutamatergic neurons. Thus, ILDs can potentially impair the responses of distinct pools of GABA_BRs, producing complex effects on the hippocampal network. The present results show that GABA_BRs on glutamatergic terminals, in contrast to those on GABAergic terminals, are not impaired following high potassium exposure.

We have previously shown that bath application of saturating concentration of baclofen does not impair postsynaptic GABA_BRs on pyramidal neurons (Tosetti *et al.*, 2004). Considering the low number of ILDs used to induce such functional impairment, we think unlikely that the postsynaptic GABA_BR-mediated control of excitability was affected in the present study. However, a recent study (Wetherington & Lambert, 2002) reported that both pre and postsynaptic GABA_BRs on glutamatergic neurons desensitised when exposed to GABA agonists for longer periods (2 to 96 hr). It is thus possible that a fully developed status epilepticus, allowing GABA_BRs to be exposed longer and to higher concentration of endogenous GABA, could affect additional pools of GABA_BR-mediated responses. We however predict that impairing the postsynaptic GABA_BR-mediated inhibition would contribute to aggravate hyperexcitability and favour the emergence of pathological activity.

Impairment of GABA auto-inhibition as a possible ictogenic mechanism

Our data demonstrate that ILDs induce the loss of GABA_BR-mediated control of GABA release in the neonatal hippocampus, thus triggering network hyperexcitability and the appearance of spontaneous pathological discharges. Among the many seizures-induced alterations that characterise

the epileptic tissue, some can underlie or facilitate the emergence of epileptiform activity while others can be seen as compensatory mechanisms to prevent seizures aggravation (Morimoto *et al.*, 2004). Whether GABA_BR activation is protective or pro-convulsive is still controversial, with effects varying greatly depending on the age, brain region and type of epileptiform activity (Engel, Jr., 1995; Bettler *et al.*, 1998; Sutor & Luhmann, 1998; Motalli *et al.*, 1999; Tosetti *et al.*, 2004; Bettler *et al.*, 2004). The protective role of GABA_BRs during development has been confirmed by the severe epileptic phenotype of GABA_{B1}-deficient mice (Prosser *et al.*, 2001; Schuler *et al.*, 2001). In the adult, however, the activation of GABA_BRs is generally considered pro-convulsive because it limits the inhibitory action of GABA_ARs (Engel, Jr., 1995). It may thus appear contradictory to evoke a loss of presynaptic inhibition of GABA release as a pro-convulsive effect. It should be considered, though, that in both adult (Kohling *et al.*, 2000; Fujiwara-Tsukamoto *et al.*, 2003) and neonatal rat hippocampus (Dzhala & Staley, 2003), GABA_A receptor activation contributes to the emergence of epileptiform activity. In the neonates, GABA_A receptors are excitatory in physiological conditions (Cherubini *et al.*, 1991). In the adult system, GABA_A receptors are usually inhibitory, but the intracellular accumulation of Cl⁻ ions during sustained GABA_AR activation (i.e. ILDs) can switch GABA_AR-mediated synaptic responses from hyperpolarising to depolarising, favouring network hyperexcitability (Staley *et al.*, 1995; Kaila *et al.*, 1997). Thus, the impairment of GABA auto-inhibition is likely to favour the emergence of pathological activity in both immature and adult animals. Several plastic changes occurring after seizures facilitate the onset of chronic epileptic activity (Morimoto *et al.*, 2004). Our data provide, however, the first evidence that GABA_BRs may contribute to post-seizure plasticity and remodelling.

It is possible that seizure-induced loss of GABA auto-inhibition contributes to the long-term hyperexcitability observed in epileptic tissue. Several studies reported a functional loss of GABA_BR-mediated inhibition (Mangan & Lothman, 1996; Haas *et al.*, 1996; Chandler *et al.*, 2003; Wu & Leung, 1997) and a decrease in the number of GABA_BRs after intense epileptic activity in adult models of chronic epilepsy (Chandler *et al.*, 2003; Straessle *et al.*, 2003; Kokaia & Kokaia, 2001). However, the mechanisms underlying the long-term impairment of GABA_BR-mediated inhibition are not known.

Our data raise the interesting possibility that seizure-induced GABA release contributes to the reduced number of GABA_BRs observed in chronic epileptic tissue. Indeed, the activity-dependent impairment of most G-protein-coupled receptors is mediated by phosphorylation followed by receptor internalisation (Tsao & von Zastrow, 2000). It is tempting to speculate that a prolonged GABA_BR activation, initially induced by ictal activity and later maintained by pathological discharges, leads to receptor internalisation and the long-term decrease in the number of functional GABA_BRs.

Our present results indicate that the functional loss of GABA auto-inhibition is a consequence of ILDs. However, a sustained release of endogenous GABA always precedes the actual development of the ILD, as well documented in Fig. 1A. It is possible, although speculative, that functional impairment of GABA_BR-mediated GABA auto-inhibition occurs during such release. If this is the case, the reduced control of GABA release might not be simply a consequence of ILDs, but could play a role in the generation of pathological discharges.

Conclusion

In the neonatal hippocampus, during physiological activity, i.e. GDPs, GABA_BR signalling is recruited to limit the release of excitatory GABA. Here we show that, during epileptiform activity, an excessive release of GABA impairs such protective mechanism, leading to aggravation of hyperexcitability and appearance of spontaneous pathological discharges. The ILD-dependent impairment of GABA auto-inhibition may thus contribute to the pro-convulsive plastic changes that characterise chronic epileptic tissues.

REFERENCES

- Andersen, K. E., Braestrup, C., Gronwald, F. C., Jorgensen, A. S., Nielsen, E. B., Sonnewald, U., Sorensen, P. O., Suzdak, P. D., & Knutsen, L. J. (1993). The synthesis of novel GABA uptake inhibitors. 1. Elucidation of the structure-activity studies leading to the choice of (R)-1-[4,4-bis(3-methyl-2-thienyl)-3-butenyl]-3-piperidinecarboxylic acid (tiagabine) as an anticonvulsant drug candidate. *J.Med.Chem.* **36**, 1716-1725.
- Ben-Ari, Y., Cherubini, E., Corradetti, R., & Gaiarsa, J.-L. (1989). Giant synaptic potentials in immature rat CA3 hippocampal neurones. *Journal of Physiology* **416**, 303-325.
- Ben-Ari, Y. & Gho, M. (1988). Long-lasting modification of the synaptic properties of rat CA3 hippocampal neurones induced by kainic acid. *Journal of Physiology* **404**, 365-384.
- Ben-Ari, Y., Khazipov, R., Leinekugel, X., Caillard, O., & Gaiarsa, J.-L. (1997). GABAA, NMDA and AMPA receptors: a developmentally regulated 'ménage a trois'. *Trends in Neurosciences* **20**, 523-529.
- Bender, R. A., Soleymani, S. V., Brewster, A. L., Nguyen, S. T., Beck, H., Mathern, G. W., & Baram, T. Z. (2003). Enhanced expression of a specific hyperpolarization-activated cyclic nucleotide-gated cation channel (HCN) in surviving dentate gyrus granule cells of human and experimental epileptic hippocampus. *Journal of Neuroscience* **23**, 6826-6836.
- Bettler, B., Kaupmann, K., & Bowery, N. (1998). GABAB receptors: drugs meet clones. *Current Opinion in Neurobiology* **8**, 345-350.

Bettler, B., Kaupmann, K., Mosbacher, J., & Gassmann, M. (2004). Molecular structure and physiological functions of GABA(B) receptors. *Physiol Rev.* **84**, 835-867.

Buhl, E. H., Otis, T. S., & Mody, I. (1996). Zinc-induced collapse of augmented inhibition by GABA in a temporal lobe epilepsy model. *Science* **271**, 369-373.

Carman, C. V. & Benovic, J. L. (1998). G-protein-coupled receptors: turn-ons and turn-offs. *Curr.Opin.Neurobiol.* **8**, 335-344.

Chandler, K. E., Princivalle, A. P., Fabian-Fine, R., Bowery, N. G., Kullmann, D. M., & Walker, M. C. (2003). Plasticity of GABA(B) receptor-mediated heterosynaptic interactions at mossy fibers after status epilepticus. *Journal of Neuroscience* **23**, 11382-11391.

Chen, K., Baram, T. Z., & Soltesz, I. (1999). Febrile seizures in the developing brain result in persistent modification of neuronal excitability in limbic circuits [see comments]. *Nat.Med.* **5**, 888-894.

Cherubini, E., Gaiarsa, J.-L., & Ben-Ari, Y. (1991). GABA: an excitatory transmitter in early postnatal life. *Trends in Neurosciences* **14**, 515-519.

Couve, A., Thomas, P., Calver, A. R., Hirst, W. D., Pangalos, M. N., Walsh, F. S., Smart, T. G., & Moss, S. J. (2002). Cyclic AMP-dependent protein kinase phosphorylation facilitates GABA(B) receptor-effector coupling. *Nat.Neurosci.* **5**, 415-424.

Davies, C. H., Davies, S. N., & Collingridge, G. L. (1990). Paired-pulse depression of monosynaptic GABA-mediated inhibitory postsynaptic responses in rat hippocampus. *Journal of Physiology* **424**, 513-531.

Dzhala, V. I. & Staley, K. J. (2003). Excitatory actions of endogenously released GABA contribute to initiation of ictal epileptiform activity in the developing hippocampus. *Journal of Neuroscience* **23**, 1840-1846.

Engel, J., Jr. (1995). Inhibitory mechanisms of epileptic seizure generation. *Adv.Neurol.* **67**, 157-171.

Ferguson, S. S., Zhang, J., Barak, L. S., & Caron, M. G. (1998). Molecular mechanisms of G protein-coupled receptor desensitization and resensitization. *Life Sci.* **62**, 1561-1565.

Freund, T. F., Gulyas, A. I., Acsady, L., Gorcs, T., & Toth, K. (1990). Serotonergic control of the hippocampus via local inhibitory interneurons. *Proc.Natl.Acad.Sci.U.S.A.* **87**, 8501-8505.

Fujiwara-Tsukamoto, Y., Isomura, Y., Nambu, A., & Takada, M. (2003). Excitatory GABA input directly drives seizure-like rhythmic synchronization in mature hippocampal CA1 pyramidal cells. *Neuroscience* **119**, 265-275.

Haas, K. Z., Sperber, E. F., Moshé, S. L., & Stanton, P. K. (1996). Kainic acid-induced seizures enhance dentate gyrus inhibition by downregulation of GABAB receptors. *Journal of Neuroscience* **16**, 4250-4260.

Holmes, G. L. & Ben-Ari, Y. (1998). Seizures in the developing brain: Perhaps not so begin after all. *Neuron* **21**, 1231-1234.

Holmes, G. L., Gaiarsa, J.-L., Chevassus-Au-Louis, N., & Ben-Ari, Y. (1998). Consequences of neonatal seizures in the rat: morphological and behavioral effects. *Annals of Neurology* **44**, 845-857.

Jensen, M. S. & Yaari, Y. (1997). Role of intrinsic burst firing, potassium accumulation, and electrical coupling in the elevated potassium model of hippocampal epilepsy. *Journal of Neurophysiology* **77**, 1224-1233.

Kaila, K., Lamsa, K., Smirnov, S., Taira, T., & Voipio, J. (1997). Long-lasting GABA-mediated depolarization evoked by high-frequency stimulation in pyramidal neurons of rat hippocampal slice is attributable to a network-driven, bicarbonate-dependent K⁺ transient. *Journal of Neuroscience* **17**, 7662-7672.

Khalilov, I., Esclapez, M., Medina, I., Aggoun, D., Lamsa, K., Leinekugel, X., Khazipov, R., & Ben-Ari, Y. (1997). A novel in vitro preparation: the intact hippocampal formation. *Neuron* **19**, 743-749.

Khalilov, I., Holmes, G. L., & Ben Ari, Y. (2003). In vitro formation of a secondary epileptogenic mirror focus by interhippocampal propagation of seizures. *Nat.Neurosci.* **6**, 1079-1085.

Khazipov, R., Leinekugel, X., Khalilov, I., Gaïarsa, J.-L., & Ben-Ari, Y. (1997). Synchronization of GABAergic interneuronal network in CA3 subfield of neonatal rat hippocampal slices. *Journal of Physiology* **498**, 763-772.

Kohling, R., Vreugdenhil, M., Bracci, E., & Jefferys, J. G. (2000). Ictal epileptiform activity is facilitated by hippocampal GABAA receptor-mediated oscillations. *Journal of Neuroscience* **20**, 6820-6829.

Kokaia, Z. & Kokaia, M. (2001). Changes in GABA(B) receptor immunoreactivity after recurrent seizures in rats. *Neuroscience Letters* **315**, 85-88.

Mangan, P. S. & Lothman, E. W. (1996). Profound disturbances of pre- and postsynaptic GABAB-receptor-mediated processes in region CA1 in a chronic model of temporal lobe epilepsy. *Journal of Neurophysiology* **76**, 1282-1296.

McLean, H. A., Caillard, O., Khazipov, R., Ben-Ari, Y., & Gañarsa, J.-L. (1996). Spontaneous release of GABA activates GABAB receptors and controls network activity in the neonatal rat hippocampus. *Journal of Neurophysiology* **76**, 1036-1046.

Morimoto, K., Fahnestock, M., & Racine, R. J. (2004). Kindling and status epilepticus models of epilepsy: rewiring the brain. *Prog. Neurobiol.* **73**, 1-60.

Motalli, R., Louvel, J., Tancredi, V., Kurcewicz, I., Wan-Chow-Wah, D., Pumain, R., & Avoli, M. (1999). GABAB receptor activation promotes seizure activity in the juvenile rat hippocampus. *Journal of Neurophysiology* **82**, 647.

Prosser, H. M., Gill, C. H., Hirst, W. D., Grau, E., Robbins, M., Calver, A., Soffin, E. M., Farmer, C. E., Lanneau, C., Gray, J., Schenck, E., Warmerdam, B. S., Clapham, C., Reavill, C., Rogers, D. C., Stean, T., Upton, N., Humphreys, K., Randall, A., Geppert, M., Davies, C. H., & Pangalos, M. N. (2001). Epileptogenesis and enhanced prepulse inhibition in GABA(B1)-deficient mice. *Mol. Cell Neurosci.* **17**, 1059-1070.

Roepstorff, A. & Lambert, J. D. C. (1994). Factors Contributing to the Decay of the Stimulus-Evoked IPSC in Rat Hippocampal CA1 Neurons. *Journal of Neurophysiology* **72**, 2911-2926.

Rutecki, P. A., Lebeda, F. J., & Johnston, D. (1985). Epileptiform activity induced by changes in extracellular potassium in hippocampus. *Journal of Neurophysiology* **54**, 1363-1374.

Schuler, V., Luscher, C., Blanchet, C., Klix, N., Sansig, G., Klebs, K., Schmutz, M., Heid, J., Gentry, C., Urban, L., Fox, A., Spooren, W., Jaton, A. L., Vigouret, J., Pozza, M., Kelly, P. H., Mosbacher, J., Froestl, W., Kaslin, E., Korn, R., Bischoff, S., Kaupmann, K., van der, P. H., & Bettler, B. (2001). Epilepsy, hyperalgesia, impaired memory, and loss of pre- and postsynaptic GABA(B) responses in mice lacking GABA(B1). *Neuron* **31**, 47-58.

Staley, K. J., Soldo, B. L., & Proctor, W. R. (1995). Ionic mechanisms of neuronal excitation by inhibitory GABAA receptors. *Science* **269**, 977-981.

Straessle, A., Loup, F., Arabadzisz, D., Ohning, G. V., & Fritschy, J. M. (2003). Rapid and long-term alterations of hippocampal GABAB receptors in a mouse model of temporal lobe epilepsy. *European Journal of Neuroscience* **18**, 2213-2226.

Sutor, B. & Luhmann, H. J. (1998). Involvement of GABAB receptors in convulsant-induced epileptiform activity in rat neocortex *in vitro*. *European Journal of Neuroscience* **10**, 3417-3427.

Tosetti, P., Bakels, R., Colin-Le Brun, I., Ferrand, N., Gáñarsa, J. L., & Caillard, O. (2004). Acute desensitization of presynaptic GABAB-mediated inhibition and induction of epileptiform discharges in the neonatal rat hippocampus. *European Journal of Neuroscience* **19**, 3227-3234.

Tsao, P. & von Zastrow, M. (2000). Down regulation of G protein-coupled receptors. *Current Opinion in Neurobiology* **10**, 365-369.

Velazquez, J. L. & Carlen, P. L. (1999). Synchronization of GABAergic interneuronal networks during seizure-like activity in the rat horizontal hippocampal slice. *European Journal of Neuroscience* **11**, 4110-4118.

Villeneuve, N., Ben-Ari, Y., Holmes, G. L., & Gaiarsa, J.-L. (2000). Neonatal seizures induced persistent changes in intrinsic properties of CA1 rat hippocampal cells. *Annals of Neurology* **47**, 729-738.

Wetherington, J. P. & Lambert, N. A. (2002). Differential desensitization of responses mediated by presynaptic and postsynaptic A1 adenosine receptors. *Journal of Neuroscience* **22**, 1248-1255.

Wu, C. & Leung, L. S. (1997). Partial hippocampal kindling decreases efficacy of presynaptic GABAB autoreceptors in CA1. *Journal of Neuroscience* **17**, 9261-9269.

Acknowledgements

This work was supported by EMBO and Marie Curie fellowships to P.T. and by the French National Institute for Medical Research (INSERM). We wish to thank Drs. Y. Ben-Ari and C. Bernard for critical reading of the manuscript. We also thank V. Rasetti, A. Sutter and A. Martinez (Novartis Pharma AG) for the gift of the GABA_B antagonist CGP35348.

FIGURE LEGENDS

Figure 1: Induction and quantification of functional impairment of GABA auto-inhibition.

A. Paired whole-cell (*top trace*) and field (*middle trace*) recordings of an intact hippocampal formation (IHF) showing the GABAergic current underlying a K-induced ictal-like discharge (ILD). The GABAergic current was isolated using a combination of voltage (+10 mV) and ionic conditions (CsGluconate patch solution). Note the continuous release of endogenous GABA over several seconds. The tonic, tonico-clonic and clonic phases of the ILD are shown at an expanded time scale (*bottom traces*). B. Left: Whole-cell GABA_A currents from a CA3 pyramidal neuron showing paired-pulse depression of GABA release (PPD) before (*control*), during (*during CGP35348*, grey trace) and after (*wash*) application of 500 μ M CGP35348 in hippocampal slices. Stimulating protocol is shown above traces. Right: average PPD before (*white bar*), during (*grey bar*) and after (*black bar*) application of 500 μ M CGP35348. Note that PPD is strongly reduced when GABA_B receptors are blocked. GABA_A currents were isolated in the presence of 10 μ M CNQX and 40 μ M APV C. Left: glutamatergic field EPSPs from CA3 *stratum radiatum* showing heterosynaptic depression of glutamate release before (*control*), during (*during CGP 35348*, grey trace) and after (*wash*) application of 500 μ M CGP35348 in hippocampal slices. Stimulating protocol is shown above traces. Right: average heterosynaptic depression before (*white bar*), during (*dark grey bar*) and after (*black bar*) application of 500 μ M CGP35348.

Asterisks indicate statistical difference with $P < 0.005$.

Figure 2: Selective impairment of GABA auto-inhibition by baclofen

A. Left: Whole-cell recordings showing PPD before (*control*), during (*during baclofen*) and after (*after baclofen*, dark grey trace) application of 100 μ M baclofen in hippocampal slices. Right: average PPD before (*white bar*), during (*black bar*) and after (*dark grey bar*) a 5 min application of 100 μ M baclofen. Note that PPD is lost following baclofen-induced functional impairment of the GABA_B-mediated response. GABA_A currents were isolated in the presence of 10 μ M CNQX and 40 μ M APV

B. Left: Field recordings from CA3 *stratum radiatum* showing heterosynaptic depression of glutamate release before (*control*), during (*during baclofen*) and after (*after baclofen*, grey trace) application of 100 μ M baclofen in hippocampal slices. Right: average heterosynaptic depression before (*white bar*), during (*black bar*) and after (*grey bar*) a 5 min application of 100 μ M baclofen. Note that no functional loss of heterosynaptic depression could be recorded after baclofen treatment.

Asterisks indicate statistical difference with $P < 0.005$.

Figure 3: Selective impairment of GABA auto-inhibition by endogenous GABA

A: Left: Traces show PPD before (*control*) and after (*after high K*) high K-induced ictal-like discharges (ILDs). GABA_A currents were isolated in the presence of 10 μ M CNQX and 40 μ M APV. Right: Average PPD before (*control*) and after ILDs (*after high K*). PPD is significantly reduced after ILDs. B. Left: Field recordings from CA3 *stratum radiatum* showing heterosynaptic depression of glutamate release before (*control*), and after ILDs (*after high K*) in hippocampal slices. Right: average heterosynaptic depression before (*white bar*), and after (*grey bar*) high K treatment. No functional loss of heterosynaptic depression could be recorded after K-induced ILDs.

Asterisks indicate statistical difference with $P < 0.005$.

Figure 4: Sustained release of endogenous GABA impairs GABA auto-inhibition

A. Left: Whole-cell recordings of CA3 pyramidal neuron showing PPD before (*control*) and after 10 mM K + 500 μ M CGP35348 treatment. Preventing GABA_B receptor activation during ILDs avoided PPD loss. Right: Average PPD before (*control*) and after 10 mM K+500 μ M CGP35348 treatment. GABA_A currents were pharmacologically isolated using 10 μ M CNQX+40 μ M APV. B. Whole-cell recordings from two CA3 pyramidal neurons in hippocampal slices. ILDs are present in 10 mM K (*High K*) but not in 10 mM K + 10 μ M CNQX + 40 μ M APV (*High K+CNQX+APV*), where only short, GDP-like GABAergic discharges can be observed. Holding potential: -65 mV. C. Average

PPD before (*control*) and after 10 mM K + 10 μ M CNQX + 40 μ M APV (*High K+CNQX+APV*). Asterisks indicate statistical difference with $P < 0.005$.

Figure 5: Impairing GABA auto-inhibition increases GDP width in hippocampal slices.

A. Voltage-clamp recordings showing GDP currents before (*Control*) and after the indicated treatments. Each trace is the mean of 15 events. $V_h = -65$ mV. B. Cumulative distribution of GDP duration before (*Control*) and after the indicated treatments. Note the significant increase in GDP duration following high K treatment. All recordings are from CA3 pyramidal neurons in acute hippocampal slices.

Figure 6: Impairing GABA auto-inhibition induces the appearance of abnormal discharges in the intact hippocampus.

A. Distribution of normal (*GDPs*) and abnormal events recorded from the intact hippocampal formation (IHF) in the presence (*after high K*) and absence (*after high K+CGP35348*) of GABA_BR activation. B. Extracellular (*Field*) and intracellular (*Whole cell*) recordings of GDPs and abnormal events (*short oscillations, long oscillations, interictal-like*) present in the IHF after high K treatment. Glutamatergic and GABAergic currents were isolated at the indicated potentials by means of a CsGluconate patch solution. The field electrode was positioned in the *stratum radiatum*. Whole cell currents were recorded from CA3 pyramidal neurons in the voltage clamp configuration.






Figure 7: Properties of abnormal events.





Frequency distributions of abnormal event properties (duration, amplitude and shape) in the presence (*after high K*) and absence (*after high K+CGP35348*) of sustained GABA_BR activation. Event amplitude and duration were normalised to control GDP values. Shape of events was expressed as number of phases, where 1 phase corresponded to 1 slope change in the trace. Note that the only values that do not significantly differ in the presence or absence of GABA_BR activation perfectly

describe the properties of interictal-like discharges. Asterisks indicate statistical difference with $P < 0.005$.



Figure 8: Partial pharmacological block of GABA_BRs induces short and long oscillations.

A. Extracellular (*Field*) and intracellular (*Whole cell*) recordings of GDPs, long and short oscillations present in an IHF during a non-saturating application of CGP35348 (100 μ M). Recording conditions were the same as in Fig. 4B. B. Frequency distributions of normal (*GDPs*) and abnormal events recorded from an IHF during 100 μ M CGP35348 (*white bars*), and after high K treatment (*black bars*). Note that the distributions are not significantly different, with the exception of interictal-like discharges, which are absent during CGP treatment.



A**IHF: 8.5 mM K⁺****B****C**





A**B**



A





B




A**B****C**


A**B**

A**B**


DURATION



SHAPE



AMPLITUDE





ADURING 100 μ M CGP35348**B**

Table 1. Properties of pathological events occurring after ILDs in the presence (*after high K*) and absence (*after high K + CGP35348*) of desensitisation of GABA auto-inhibition. Table values represent the % of total events having the specified duration, amplitude or shape. Amplitude and duration bins are expressed as % of control GDP values. Asterisks indicate a statistically significant reduction ($P<0.05$).

	after high K (%)	after high K + CGP (%)
duration		
<100%	6.0±4.1	3.9±2.7
100-200%	16.1±5.5	16.9±7.1
200-300%	31.7±8.0	5.2±3.8 *
300-400%	8.7±4.7	1.6±1.6 *
400-1000%	4.9±3.0	1.9±1.9 *
>1000%	10.7±0.1	—
amplitude		
<100%	4.6±2.8	—
100-200%	18.0±6.6	3.6±2.2 *
200-300%	8.3±4.7	1.5±1.5 *
300-500%	17.4±6.2	17.6±6.8
500-1000%	14.6±5.2	3.1±2.1 *
>1000%	13.0±6.0	3.6±3.6 *
phases		
2	1.6±1.2	1.6±1.6
3	18.9±6.8	19.2±7.0
4	19.3±6.2	6.6±3.8 *
5	10.6±4.6	2.1±2.1 *
6	7.0±3.9	—
7 to 15	10.8±4.2	—
>15	13.0±5.3	—